# HOMOLOGY THEORY Notes & Exercises from my Independent Study

(OR: If I could save Klein in a bottle  $\checkmark$ )

## FOREST KOBAYASHI

Department of Mathematics Harvey Mudd College

 $\begin{array}{l} {\rm Supervised} \ {\rm By} \\ {\rm Francis} \ {\rm Su}^1 \end{array}$ 



<sup>&</sup>lt;sup>1</sup>Department of Mathematics, Harvey Mudd College

# Contents

Introdu	action	v
15 Hon	nological Prereqs	1
15.1	Manifolds	1
15.2	Simplicial Complexes	1
15.3	Simplicial Maps and PL Homeomorphisms	5
15.4	Simplicial Approximation	5
<b>16</b> Z/2Z	Z Homology	1
16.1	Intro	11
16.2	Chains, Cycles, Boundaries, and the Homology Groups	11
16.3	Induced Homomorphisms and Invariance	23
16.4	The Mayer-Vietoris Theorem	27
$18 \mathbb{Z} H$	omology	37
18.1	Orientation and Z-Homology	37
18.2	Relative Simplicial Homology	39
18.3	Useful Exact Sequences	46
18.4	The Degree of a Map	46
18.5	Lefschetz	47
19 Som	e Homological Algebra 4	19
19.1	Chain complexes	49
20 Rot	man	51
20.1	A sketch of the Brouwer Fixed Point Theorem	51
21 App	pendix	53
21.1	List of Definitions	53

### Introduction

#### What's this?

This document is a compendium of notes, exercises, and other miscellany from my independent study in Homology Theory. For this, I am working through the second half of *Topology Through Inquiry* by Michael Starbird and Francis Su (i.e., chapters 11-20), under supervision from Prof. Su himself. Rough topic coverage should be discernable from the table of contents, as I've tried to name each section identically to the corresponding title in the book.

#### Notation

Most notation I use is fairly standard. Here's a (by no means exhaustive) list of some stuff I do.

- "WTS" stands for "want to show," s.t. for "such that." WLOG, as usual, is without loss of generality.
- End-of-proof things: is QED for exercises and theorems. □ is used in recursive proofs (e.g., proving a Lemma within a theorem proof). If doing a proof with casework, ✓ will be used to denote the end of each case.
- $(\Rightarrow \Leftarrow)$  means contradiction
- $\mathcal{T}(U)$  will denote the topology of a topological space U.
- $\mathcal{P}(A)$  is the powerset of A. I don't like using  $2^A$ .
- $\rightarrow$  denotes surjection.
- $\hookrightarrow$  denotes injection.
- Thus,  $\hookrightarrow$  denotes bijection.
- Important: I use f→(A) for the image of A under f, and f←(B) for the inverse image of B under f.
- $\sim$  and  $\equiv$  are used for equivalence relations.  $\cong$  is used to denote homeomorphism and isomorphism of groups.  $\simeq$  is for Homotopy equivalence.
- $\epsilon$  is for trivial elements (e.g., the trivial path), while  $\epsilon$  is for small positive quantities.
- $\overline{U}$  denotes the closure of  $U, U^{\circ}$  is the interior of U.
- $A^c$  is A complement.
- $\{v_0 \cdots v_k\}$  denotes a simplex on k + 1 vertices (that is, a k-simplex).  $\{v_0 \cdots \hat{v_i} \cdots v_k\}$  is the same simplex with the  $i^{\text{th}}$  vertex deleted.
- $[n] = \{i \mid i = 0, 1, \dots, n\}.$

## 15. Manifolds, Simplexes Complexes, and Triangulability: Building Blocks

#### 15.1 Manifolds

We define some basic Euclidean sets for use in homeomorphisms.

**Definition 15.1.1** (*n*-cube). The *n*-dimensional cube, denoted  $\mathbb{D}^n$ , is defined as

$$\mathbb{D}^n = \{ (x_1, \dots, x_n) \in \mathbb{R}^n \mid 0 \le x_i \le 1 \text{ for } i = 1, \dots, n \}$$
$$= \overbrace{[0,1] \times [0,1] \times \dots \times [0,1]}_{n \text{ times}} \subset \mathbb{R}^n.$$

**Definition 15.1.2** (*n*-ball). The standard *n*-ball, denoted  $B^n$ , is

$$B^{n} = \{ (x_{1}, \dots, x_{n}) \in \mathbb{R}^{n} \mid x_{1}^{2} + \dots + x_{n}^{2} \le 1 \}.$$

**Definition 15.1.3** (*n*-sphere). The standard *n*-sphere, denoted  $\mathbb{S}^n$ , is

$$\mathbb{S}^{n} = \{ (x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid x_0^2 + \dots + x_n^2 = 1 \}.$$

note that here, our indices start at 0.

**Definition 15.1.4** (*n*-manifold). An *n*-dimensional manifold or *n*-manifold is a separable metric space M such that  $\forall p \in M, \exists U \in \mathcal{T}(M)$  s.t.  $p \in U$  and  $U \cong V \subset \mathbb{R}^n$ .

**15.8.** If M is an *n*-manifold and U is an open subset of M, then U is also an *n*-manifold.

**15.9.** If M is an m-manifold and N is an n-manifold, then  $M \times N$  is an (m+n)-manifold.

**15.10.** Let  $M^n$  be an *n*-dimensional manifold with boundary. Then  $\partial M^n$  is an (n-1)-manifold.

## 15.2 Simplicial Complexes

**Definition 15.2.1** (Affine Independence). Let  $X = \{v_0, \ldots, v_k\} \subset \mathbb{R}^n$ . We say X is affinely independent if  $\{v_1 - v_i, \ldots, v_k - v_i\}$  is linearly independent for all  $v_i$ .

**Example 15.2.1.**  $X = \{(0,1), (-\sqrt{3}/2, -1/2), (\sqrt{3}/2, -1/2)\}$  is affinely independent.

**Definition 15.2.2** (Convex combination). A *convex combination* of  $v_0, \ldots, v_k$  is a linear combination of these points whose coefficients are nonnegative and sum to 1.

**Definition 15.2.3** (k-simplex). A k-simplex is the set of all convex combinations of k + 1 affinely independent points in  $\mathbb{R}^n$ . For affinely independent points  $v_0, \ldots, v_k$  in  $\mathbb{R}^n$ ,  $\{v_0 \cdots v_k\}$  denotes the

k-simplex

$$\left\{\lambda_0 v_0 + \lambda_1 v_1 + \dots + \lambda_k v_k \mid \forall i = 0, 1, \dots, k; \ 0 \le \lambda_i \le 1 \text{ and } \sum_{i=0}^k \lambda_i = 1\right\}$$

each  $v_i$  is called a *vertex* of  $\{v_0 \cdots v_k\}$ . Any point x in the k-simplex is specified uniquely by the k+1 coefficients  $(\lambda_i)$ ; these coefficients are called the *barycentric coordinates of x*. The *barycentric coordinate of x with respect to vertex v<sub>i</sub>* is the coefficient  $\lambda_i$ .

**Definition 15.2.4** (Face and dimension). Any simplex  $\tau$  whose vertices are a nonempty subset of the vertices of a k-simplex  $\sigma$  is called a *face* of  $\sigma$ . If the number of vertices is i + 1, then  $\tau$  has *dimension* i and is called an *i*-face of  $\sigma$  and  $\tau$  has *codimension* k - i, the number of dimensions below the top dimension.

**Notational Note:** if  $\sigma = \{v_0 \cdots v_k\}$ , the (k-1)-dimensional face of  $\sigma$  obtained by deleting the vertex  $v_i$  from the list of vertices of  $\sigma$  is denoted by  $\{v_0 \cdots \hat{v_i} \cdots v_k\}$ .

**15.11.** Show that if  $\sigma$  is a simplex and  $\tau$  is one of its faces, then  $\tau \subset \sigma$ .

Solution. This is fairly trivial, so we offer just a sketch. Suppose  $\mathbf{v} \in \tau$ . Then write  $\mathbf{v}$  as an element of  $\sigma$  by taking  $\lambda_i = 0$  for all those  $v_i \notin \tau$ .

**Definition 15.2.5** (Simplicial complex). A simplicial complex K (in  $\mathbb{R}^n$ ) is a collection of simplicies in  $\mathbb{R}^n$  satisfying the following conditions.

- 1. If a simplex  $\sigma$  is in K, then each face of  $\sigma$  is also in K.
- 2. Any two simplices in K are either disjoint or their intersection is a face of each.

**15.13.** Exhibit a collection of simplices that satisfies condition (1) but not condition (2) in the definition of a simplicial complex.

*Solution.* Consider the following diagram, where the interior of each simplex is taken to be in the complex.



Figure 15.1: An unfortunate collision

Note that to fix this sorry situation, we can't just add two vertices at the points of intersections of the lines above (then the intersection of the resulting simplex with the two shone above would be non-trivial, but still not a face of the larger ones). We'd actually need something much more complicated.



Figure 15.2: Constructing a resolution



Figure 15.3: The completed resolution

**Definition 15.2.6** (Underlying space). The *underlying space* |K| of a simplicial complex K is the set

$$|K| = \bigcup_{\sigma \in K} \sigma,$$

the union of all simplices in K, with a topology consisting of sets whose intersection with each simplex  $\sigma \in K$  is open in  $\sigma$ . For finite simplicial complexes, this topology is the topology inherited as a subspace of  $\mathbb{R}^n$ .

**15.14.** Let K be the following simplicial complex:

(Omitted because it takes a long time to TeX out)

draw K and its underlying space.

Solution.



Figure 15.4: K (left) and its underlying space (right).

**Definition 15.2.7** (Triangulable). A topological space X is said to be *triangulable* if it is homeomorphic to the underlying space of a simplicial complex K. In that case, we say K is a *triangulation* of X.

**15.15.** Show that the space shown in Figure 15.2 (not included here) is triangulable by exhibiting a simplicial complex whose underlying space it is homeomorphic to.

Solution.



Figure 15.5: Such a simplicial complex. Note, the left triangle is unfilled.

**15.6.** For each  $n \in \mathbb{N}$ ,  $\mathbb{S}^n$  is triangulable.

*Proof.* We proceed by induction.

**Base Case:** Note that  $S^0$  is trivially triangulable by taking  $K = \{\{v_0\}, \{v_2\}\}$ .

**Inductive Hypothesis** Suppose that for  $k \in \mathbb{N} \cup \{0\}$ ,  $\mathbb{S}^k$  is triangulable by a simplicial complex K.

**Inductive Step:** Take  $v_{k+1} \in \mathbb{R}^{k+1}$  such that  $v_{k+1} \in (\operatorname{span}(K))^{\perp}$ . Then

Last Updated April 19, 2019

This proof is unfinished. Hey, future Forest — you should return to this later!

## 15.3 Simplicial Maps and PL Homeomorphisms

We now define structure-preserving maps bewteen simplicial concepts.

**Definition 15.3.1** (Simplicial map). Let X, Y be topological spaces. A function  $f : X \to Y$  is called a *simplicial map* iff there exist simplicial complexes K and L such that |K| = X, |L| = Y, and f maps each simplex of K linearly onto a (possibly lower-dimensional) simplex in L.

**Definition 15.3.2** (Simplicially homeomorphic). A simplicial map f is a simplicial homomorphism iff it's a bijection; in that case, the two complexes are *simplicially homeomorphic* 

**15.17.** A simplicial map from K to L is determined by the images of the vertices of K.

Solution. Apply linearity and show the analog of images of liner combinations being uniquely determined by the action on the basis.

**15.18.** A composition of simplicial maps is a simplicial map.

Solution. Simply plug in arbitrary points and verify the properties hold.

**Definition 15.3.3** (Subdivision). Let K be a simplicial complex. Then a simplicial complex K' is a *subdivision* of K iff each simplex of K' is a subset of a simplex of K and each simplex of K is the union of finitely many simplices of K'.

**Definition 15.3.4** (Piecewise linear). If K and L are complexes, a continuous map  $f : |K| \to |L|$  is called *piecewise linear* or PL if and only if there are subdivisions K' of K and L' of L such that f is a simplicial map from K' to L'. If there exist subdivisions such that f is a simplicial homomorphism, then f is a PL homeomorphism and the spaces are PL homeomorphic.

**15.21.** A composition of PL maps is PL. A PL homeomorphism is an equivalence relation.

Solution. Let K, L, M be complexes, and let  $g : |K| \to |L|, f : |L| \to |M|$  be continuous PL maps. WTS  $h = f \circ g$  is a PL map.

Let K', L', M' be the corresponding subdivisions of K, L, and M, respectively. Then g is a simplicial map from K' to L', and f is a simplicial map from L' to M'. Then  $\forall \sigma \in K'$ ,  $g(\sigma) \in L'$ , whence  $f(g(\sigma)) \in M'$ . It follows that  $h = g \circ f$  is a simplicial map from K' to M'.

We give a sketch of the proof that PL homeomorphism is an equivalence relation. To verify reflexivity, take the identity map. Symmetry follows by inverting the simplicial homeomorphism. Transitivity follows by the above. Thus, the claim holds.

## 15.4 Simplicial Approximation

**15.23.** Let K be a complex consisting of the boundary of a triangle (three vertices and three edges) and L be an isomorphic complex. Both |K| and |L| are topologically circles. There is a continuous map that takes the circle |K| and winds it twice around the circle |L|; however, show that there is no simplicial map from K to L that winds the circle |K| twice around the circle |L|.

Solution. We offer a brief sketch. Basically, this would require each 1-simplex to map to two 1-simplices. Contradiction.

**Definition 15.4.1** (Barycenter). The *barycenter* of a k-simplex  $\{v_0 \cdots v_k\}$  in  $\mathbb{R}^n$  is the point  $\frac{1}{k+1}(v_0 + \cdots + v_k)$ .

**Definition 15.4.2** (First barycentric subdivision  $(\operatorname{sd} \sigma)$ ). Let  $\sigma$  be an *n*-simplex. The first barycentric subdivision of  $\sigma$ , denoted  $\operatorname{sd} \sigma$ , is the complex of all simplices of the form  $\{b_0 \cdots b_k\}$ , where  $b_i$  is the barycenter of a face  $\sigma^i$  of  $\sigma$  from a chain of faces of  $\sigma$ ,

$$\sigma^0 \subset \sigma^1 \subset \cdots \subset \sigma^k$$

of increasing (not necessarily consecutive) dimensions. The maximal simplices, that is, the *n*-simplices of  $sd \sigma$  each arise from a maximal sequence of faces, that is, from faces of consecutive dimensions starting with a vertex of  $\sigma$ . Thus an *n*-simplex of  $sd \sigma$  corresponds exactly to a permutation of the vertices of  $\sigma$ .

**Definition 15.4.3** (sd K). Let K be a simplicial complex. The *first barycentric subdivision* of K, denoted sd K, is the complex consisting of all the simplices in the barycentric subdivision of each simplex of K.

**Definition 15.4.4** (*m*-th barycentric subdivision). The second barycentric subdivision, denoted  $sd^2 K$ , is the first barycentric subdivision of sd K. Proceeding in this way, the *m*-th barycentric subdivision is denoted  $sd^m K$ .

Some diagrams:



Figure 15.6: The first 4 barycentric subdivisions

15.24. How many *n*-simplices are there in the first barycentric subdivision of an *n*-simplex?

Solution. A simple induction shows there are  $6^n$  *n*-simplices.

**15.25.** Convince yourself that the barycentric subdivision of a complex K is, in fact, a subdivision of K.

Solution. I'm convinced.

**15.26.** Let K be a finite simplicial complex and let  $a_n$  be the maximum among the diameters of simplices in  $sd^n K$ . Then

 $\lim_{n \to \infty} a_n = 0.$ 

Solution. First, we calculate the diameter of an *n*-simplex.

**Lemma 15.4.1.** Let  $\sigma_n$  be an n-simplex. Then the diameter of  $\sigma_n$ 

$$D = \sup_{\mathbf{x}, \mathbf{y} \in \sigma_n} \|\mathbf{x} - \mathbf{y}\|_2$$

is given by the maximum distance between vertices in the simplex:

$$D = \sup_{\mathbf{v}_i, \mathbf{v}_j} \|\mathbf{v}_i - \mathbf{v}_j\|_2$$

*Proof of Lemma:* Let  $\mathbf{x}, \mathbf{y} \in \sigma_n$  be arbitrary. It will suffice to show that  $\mathbf{y}$  is not a vertex in  $\sigma_n$ , then there exists a vertex  $\mathbf{v} \in \sigma_n$  such that  $\|\mathbf{x} - \mathbf{y}\|_2 < \|\mathbf{x} - \mathbf{v}\|_2$ .

Write  $\mathbf{y}$  as convex combinations by

$$\mathbf{y} = \sum_{i=0}^{n} \mu_i \mathbf{v}_i.$$

and observe that since  $\sum_{i=0}^{n} \mu_i = 1$ , we have

$$\mathbf{x} = \sum_{i=0}^{n} \mu_i \mathbf{x} = \mathbf{x}.$$

Then

$$\|\mathbf{x} - \mathbf{y}\|_{2} = \left\| \sum_{i=0}^{n} \mathbf{x} - \mu_{i} \mathbf{v}_{i} \right\|_{2}$$
$$= \left\| \sum_{i=0}^{n} \mu_{i} (\mathbf{x} - \mathbf{v}_{i}) \right\|_{2}$$
$$\leq \sum_{i=0}^{n} \mu_{i} \|\mathbf{x} - \mathbf{v}_{i}\|_{2}$$
$$\leq \sum_{i=0}^{n} \mu_{i} \sup_{\mathbf{v}_{i}} \|\mathbf{x} - \mathbf{v}_{i}\|_{2}$$
$$= \sup_{\mathbf{v}_{i}} \|\mathbf{x} - \mathbf{v}_{i}\|_{2}.$$

Hence, we see for arbitrary  $\mathbf{y}$ ,  $\|\mathbf{x} - \mathbf{y}\|_2 \le \sup_{\mathbf{v}_i} \|\mathbf{x} - \mathbf{v}_i\|_2$ . Now, apply the same result to  $\mathbf{x}' = \arg \max_{\mathbf{v}_i} \|\mathbf{x} - \mathbf{v}_i\|_2$  and  $\mathbf{y}' = \mathbf{x}$  to obtain

$$\|\mathbf{x} - \mathbf{y}\|_{2} \leq \sup_{\mathbf{v}_{i}} \|\mathbf{x} - \mathbf{v}_{i}\|_{2} \leq \sup_{\mathbf{v}_{j}} \left( \sup_{\mathbf{v}_{i}} \|\mathbf{v}_{j} - \mathbf{v}_{i}\| \right) = \sup_{\mathbf{v}_{i}, \mathbf{v}_{j}} \|\mathbf{v}_{j} - \mathbf{v}_{i}\|_{2}$$

as desired.

By the lemma,  $a_n$  is given by the maximal side length of a 2-simplex in  $sd^n K$ . Hence

$$0 \le a_n \le \frac{1}{2^n} \frac{2}{\sqrt{3}}$$
 This bound is incorrect. How can I fix it

and so by the squeeze theorem,

$$\lim_{n \to \infty} a_n = 0$$

as desired.

**Definition 15.4.5** (Minimal face). Let K be a simplicial complex. The minimal face of  $x \in |K|$  is the simplex of K of smallest dimension that contains x.

**Definition 15.4.6** (Star of vertex). The star of a vertex v in K, denoted St(v), is the set of all points whose minimal face contains v.

**Remark.** The definition of the star of a vertex is basically the interior of the union of all simplices containing v.

**15.27.** The star of a vertex v in a complex K is an open set of |K|, and the collection of all vertex stars covers |K|.

Solution. Let  $v \in K$  be a vertex. Let  $x \in St(v)$  be arbitrary. WTS  $\exists \varepsilon > 0$  s.t.  $B_{\epsilon}(x) \subset St(v)$ . We have the following cases:

- (1) Suppose x = v. Then taking  $\epsilon = \frac{1}{2} \inf_{\mathbf{v}_i} |v \mathbf{v}_i|_2$  we get the desired result.
- (2) Suppose  $x \neq v$ . Then taking  $v = v_0$ , write x in the barycentric coordinates

 $x = \lambda_0 v + \lambda_1 v_1 + \dots + \lambda_n v_n.$ 

Since  $x \in \operatorname{St}(v), \lambda_0 \neq 0$ .

**15.28.** If the simplex  $\sigma = \{v_0 \cdots v_k\}$  in K is the minimal face of a point  $x \in |K|$ , then

$$x \in \bigcap_{i=0}^{n} \operatorname{St}(v_0)$$

Solution.

Forest Kobayashi

## 16. Simplicial $\mathbb{Z}_2$ -Homology: Physical Algebra

#### 16.1 Intro

This chapter, we'll talk about *homology*, which captures holes in a much more satisfying way than higher homotopy groups do.

**Remark.** Although not exactly accurate, a good way to start to understand homology for a space X is to view an n-manifold in X that is not the boundary of an (n + 1)-manifold-with-boundary as capturing some geometry of X while an n-manifold that is the boundary of an (n + 1)-dimensional manifold-with-boundary is not detecting any hole or structure.

# 16.2 Chains, Cycles, Boundaries, and the Homology Groups Definition 16.2.1 (*n*-chain). An *n*-chain of K is a finite formal sum

$$\sum_{i=1}^k \sigma_i$$

of distinct n-simplices in K. Note that the dimensions of the simplices must be the same. So *chain* will mean n-chain whenever the dimension is either unimportant or understood.

**Definition 16.2.2** (*n*-chain group). The *n*-chain group of K (with coefficients in  $\mathbb{Z}/2\mathbb{Z}$ ), denoted  $C_n(K)$ , is the collection of *n*-chains in K under formal addition modulo 2. If there are no *n*-simplices in K, the *n*-chain group of K is defined to be trivial (containing the "empty" chain).

**16.1.** Check that  $C_n(K)$  is an abelian group.

Solution.

(1) 
$$\epsilon = \sum_{i \in \emptyset} \sigma_i$$

- (2) Associativity inherited from  $\cup$ .
- (3) Closure inherited from  $\cup$  over the domain given.
- (4) Existence of inverses since we're taking formal linear combinations over  $\mathbb{Z}/2\mathbb{Z}$ , then every element is its own inverse.

Finally, to see that  $C_n(K)$  is abelian, observe that + in  $C_n(K)$  inherits commutativity from  $\cup$ .

**Definition 16.2.3** ( $\mathbb{Z}/2\mathbb{Z}$  boundary of a simplex). The  $\mathbb{Z}/2\mathbb{Z}$ -boundary of an *n*-simplex  $\sigma = \{v_0 \cdots v_n\}$  is defined by

$$\partial \sigma = \sum_{i=0}^{n} \{ v_0 \cdots \widehat{v_i} \cdots v_n \}$$

the formal sum of the (n-1)-faces of  $\sigma$ .

For a 0-simplex, the  $\mathbb{Z}/2\mathbb{Z}$  boundary is defined to be  $0 \in C_{-1}(K)$ .

**Definition 16.2.4** ( $\mathbb{Z}/2\mathbb{Z}$  boundary of an *n*-chain). The  $\mathbb{Z}/2\mathbb{Z}$  boundary of an *n*-chain is the sum of the boundaries of the simplices. That is,  $\partial_n : C_n(K) \to C_{n-1}(K)$  is given by

$$\partial\left(\sum_{i=1}^k \sigma_i\right) = \sum_{i=1}^k \partial(\sigma_i)$$

16.2. Verify that  $\partial$  is a homomorphism, and use the definition to compute the  $\mathbb{Z}/2\mathbb{Z}$  boundary of  $\sigma_1 + \sigma_2$  in Figure 16.1

Solution. We want to show  $\partial$  is a homomorphism.

(a) Let  $\epsilon_n \in \mathsf{C}_n(K)$  be identity. We want to show  $\partial(\epsilon_n) = \epsilon_{n-1}$ . Taking the empty sum to be identity, we see

$$\partial(\epsilon_n) = \partial\left(\sum_{i \in \emptyset} \sigma_i\right)$$
$$= \sum_{i \in \emptyset} \partial(\sigma_i)$$
$$= \epsilon_{n-1}$$

as desired.

(b) That  $\partial$  respects addition is definitional.

We have  $\partial(\sigma_1 + \sigma_2) = e_1 + e_2 + e_4 + e_5$ .

**Definition 16.2.5** (*n*-cycle and *n*-boundary). An *n*-cycle is an *n*-chain of *K* whose boundary is zero. The set of all *n*-cycles on *K* is denoted  $Z_n(K)$ . An *n*-boundary is an *n*-chain that is the boundary of an (n + 1)-chain of *K*. The set of all *n*-boundaries is denoted  $B_n(K)$ .

**16.4.** Both  $Z_n(K)$  and  $B_n(K)$  are subgroups of  $C_n(K)$ . Moreover,

 $\partial\circ\partial=0.$ 

In other words,  $\partial_n \circ \partial_{n+1} = 0$  for each index  $n \ge 0$ . Hence,  $\mathsf{B}_n(K) \subset \mathsf{Z}_n(K)$ .

Solution. Let  $\sigma_1, \sigma_2 \in \mathsf{Z}_n(K)$ . Then by linearity of  $\partial_n$ , we have

$$\partial_n(\sigma_1 + \sigma_2) = \partial_n(\sigma_1) + \partial_n(\sigma_2)$$
  
= 0

and hence  $\mathsf{Z}_n(K) < \mathsf{C}_n(K)$ .

Now, let  $\sigma_1, \sigma_2 \in \mathsf{B}_n(K)$ . Then  $\exists \tau_1, \tau_2 \in \mathsf{Z}_{n+1}(K)$  such that  $\partial_{n+1}(\tau_1) = \sigma_1, \partial_{n+1}(\tau_2) = \sigma_2$ . Since  $\mathsf{Z}_{n+1}(K) < \mathsf{C}_{n+1}(K)$ , then  $\tau_1 + \tau_2 \in \mathsf{Z}_{n+1}(K)$ . Now, by linearity of  $\partial$ , we have

$$\partial_{n+1}(\tau_1 + \tau_2) = \partial_{n+1}(\tau_1) + \partial_{n+1}(\tau_2)$$
$$= \sigma_1 + \sigma_2$$

hence  $\mathsf{B}_n(K)$  is a subset closed under the operation, so we have  $\mathsf{B}_n(K) < \mathsf{C}_n(K)$ .

It remains to show  $\partial_n \circ \partial_{n+1} = 0$ . Let  $\sigma \in \mathsf{C}_{n+1}(K)$ . Then

$$\partial_{n+1}(\sigma) = \partial_{n+1} \left( \sum_{i \in I} \left\{ v_0^{(i)} \cdots v_{n+1}^{(i)} \right\} \right)$$
$$= \sum_{i \in I} \partial_{n+1} \left( \left\{ v_0^{(i)} \cdots v_{n+1}^{(i)} \right\} \right)$$
$$= \sum_{i \in I} \sum_{j \in [n+1]} \left\{ v_0^{(i)} \cdots \widehat{v_j^{(i)}} \cdots v_{n+1}^{(i)} \right\}$$

and so

$$\partial_n(\partial_{n+1}(\sigma)) = \sum_{i \in I} \sum_{j \in [n+1]} \partial_n \left( \left\{ v_0^{(i)} \cdots \widehat{v_j^{(i)}} \cdots v_{n+1}^{(i)} \right\} \right)$$
$$= \sum_{i \in I} \sum_{j \in [n+1]} \sum_{\substack{k \in [n+1]\\k \neq j}} \left\{ v_0^{(i)} \cdots \widehat{v_k^{(i)}} \cdots \widehat{v_j^{(i)}} \cdots v_{n+1}^{(i)} \right\}$$

hence all the terms cancel, and we're left with **0**. So  $\partial_n \circ \partial_{n+1} = 0$ , as desired.

Since every  $\sigma \in \mathsf{B}_n(K)$  is of the form  $\partial_{n+1}(\tau)$  where  $\tau \in \mathsf{C}_{n+1}(K)$ , it follows that  $\partial_n^{\rightarrow}(\mathsf{B}_n(K)) = (\partial_n \circ \partial_{n+1})(\mathsf{C}_{n+1}(K)) = 0$ . Thus  $\mathsf{B}_n(K) < \mathsf{Z}_n(K)$ .

**Definition 16.2.6** (Homologous cycles). Two *n*-cycles  $\alpha$  and  $\beta$  in *K* are *equivalent* or *homologous* iff  $\alpha - \beta = \partial(\gamma)$  for some (n + 1)-chain  $\gamma$ . In other words,  $\alpha$  and  $\beta$  are homologous iff they differ by an element of the subgroup  $\mathsf{B}_n(K)$ , denoted by

$$\alpha \sim_{\mathbb{Z}/2\mathbb{Z}} \beta.$$

The equivalence class of  $\alpha$  is denoted by enclosing it in brackets thusly:  $[\alpha]$ . For  $\mathbb{Z}/2\mathbb{Z}$  *n*-chains, observe that  $\alpha - \beta = \alpha + \beta$ . So we see that two *n*-cycles are equivalent if together they bound an (n+1)-chain.

**Definition 16.2.7** ( $n^{\text{th}}$  Homology group). The  $n^{\text{th}}$ -homology group (with coefficients in  $\mathbb{Z}/2\mathbb{Z}$ ) of a finite simplicial complex K, denoted  $\mathsf{H}_n(K)$ , is the additive group whose elements are equivalence classes of cycles under the  $\mathbb{Z}/2\mathbb{Z}$ -equivalence defined above, with  $[\alpha] + [\beta] = [\alpha + \beta]$ . I.e.,

$$\mathsf{H}_n(K) = \mathsf{Z}_n(K) / \mathsf{B}_n(K)$$

**F1.** Consider the simplicial complex given below in Figure (lol oops). Then for n = 0, 1, 2,

- (a) describe elements of  $C_n(K)$ ,
- (b) compute  $\mathsf{Z}_n(K)$ ,
- (c) compute  $\mathsf{B}_n(K)$ , and
- (d) compute  $H_n(K)$ .



Figure 16.1: Simplicial complex K

Solution. First, we redraw the simplicial complex as follows:



Figure 16.2: Simplicial complex K, straightened out

For the purposes of this problem, take angled brackets indicate span. We have

- (i) We calculate the k = 0 case.
  - (a) Elements of  $C_0(K)$  are formal linear combinations over the set  $\{v_1, v_2, \ldots, v_6\}$ . Then ١ 1 (

$$\mathsf{C}_0(K) = \left\langle v_1, v_2, v_3, v_4, v_5, v_6 \right\rangle$$

that is, collections of points in  $\mathsf{C}_0(K).$ 

(b) Let  $\sigma_1, \ldots, \sigma_k \in \mathsf{C}_0(K)$ . Then by definition,

$$\partial\left(\sum_{i=1}^{k}\sigma_{i}\right) = \sum_{i=1}^{k}\partial(\sigma_{i})$$
$$= \sum_{i=1}^{k}0$$
$$= 0$$

 $14 \ {\rm of} \ 54$ 

hence  $\mathsf{Z}_n(K) = \mathsf{C}_n(K)$ .

(c) A  $\sigma \in \mathsf{C}_0(K)$  is an *n*-boundary if  $\exists \tau \in \mathsf{C}_1(K)$  with  $\partial(\tau) = \sigma$ . Note, for any 1-dimensional face  $\{v_i v_j\} \in K$ ,

$$\partial(\{v_i v_j\}) = \{v_i \hat{v}_j\} + \{\hat{v}_i v_j\}$$
$$= \{v_i\} + \{v_j\}$$
$$= \delta_{ij}.$$

Hence, any edge formed of a pair of two distinct vertices yields a nonempty boundary. We first count all edges:



Figure 16.3: Simplicial complex K with simple edges

Since  $B_0(K)$  is a subgroup of  $C_0(K)$ , by closure under +, we see that any  $v_i + v_j$ in K such that there exists a path from  $v_i$  to  $v_j$  (when K is considered a graph) is an element of  $B_0(K)$ . In fact, we can say more:

**Claim:** Since K is connected as a graph, any even collection of vertices is in  $B_n(K)$ .

**Proof of Claim:** Suppose we have  $\sigma = \{v_{i_1}\} + \{v_{i_2}\} + \dots + \{v_{i_{2k}}\}$ , where  $k \in \mathbb{N}$ . Then for each  $j = 1, \dots, k$ , let  $\tau_j$  be a sum of edges representing a path from  $v_{i_j}$  to  $v_{i_{j+1}}$ . For example, if  $v_{i_j} = v_6$  and  $v_{i_{j+1}} = v_2$ , we could take the following approaches:





Figure 16.4: Some paths from  $v_6$  to  $v_2$ 

among others. Taking the sum of the constituent edges in each path yields a sum of 1-simplices with boundary  $v_6, v_2.^1$ 

- (d) Since  $\mathsf{B}_n(K)$  is the group of all collections of even vertices in  $\mathsf{C}_n(K)$ , we have  $\mathsf{H}_n(K) = \mathsf{C}_n(K)/\mathsf{B}_n(K) \cong \mathbb{Z}/2\mathbb{Z}$ .
- (ii) Now, we calculate the k = 1 case.
  - (a) Elements of  $C_1(K)$  are collections of linear combinations of the edges

$$\mathsf{C}_1(K) = \langle e_1, e_2, e_3, e_4, e_5, e_6, e_7, e_8, e_9 \rangle$$

(b) Elements of  $Z_1(K)$  are collections of edges such that each vertex contained in an edge in the collection has even degree. This corresponds to cyclic subgraphs of K (as well as the empty cycle), e.g.:



<sup>&</sup>lt;sup>1</sup>Justification: note that the coefficient on any given vertex when we apply  $\partial$  is the degree of the vertex in our path. Hence, only the initial and terminal vertex don't get mapped to 0.



Figure 16.5: Some cycles in K

(c) First, consider the following diagram:



Figure 16.6: Two n = 2 simplices

 $\mathbf{0}_1$  bounds  $\mathbf{0}_2$ . Since  $\partial(A_1) \cap \partial(A_2) = \emptyset$ , then the other two cycles in  $\mathsf{B}_1(K)$  are just  $\partial(A_1)$  and  $\partial(A_2)$ , respectively.

- (d)  $H_1(K) \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$  (equivalence classes have representative elements  $\mathbf{0}, \partial(A_1), \partial(A_2), \partial(A_1) + \partial(A_2)$ )
- (iii) For k = 2, we have
  - (a)  $\mathsf{C}_2(K) \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$
  - (b)  $\mathsf{Z}_2(K) \cong \mathbf{0}$

- (c)  $\mathsf{B}_2(K) \cong \mathbf{0}$
- (d) And hence  $H_2(K) \cong \mathbf{0}$ .

**16.7.** If K is a one-point space,  $H_n(K) \cong 0$  for  $n \ge 0$ , and  $H_0(K) \cong \mathbb{Z}/2\mathbb{Z}$ .

Solution. For n > 0,  $C_n(K)$  is the trivial group. Since  $Z_n(K) \leq C_n(K)$ , we thus have  $Z_n(K) \cong 0$ , and so  $H_n(K) \cong 0$ .

For the n = 0, note that  $\mathsf{Z}_0(K) = \mathsf{C}_0(K) \cong \mathbb{Z}/2\mathbb{Z}$  (every point is definitionally a 0-cycle). Since K contains no 1-simplices,  $\mathsf{B}_0(K) = \mathbf{0}$ , hence  $\mathsf{H}_0(K) \cong \mathbb{Z}/2\mathbb{Z}$ .

**Definition 16.2.8** (Acyclic). Any space with the homology groups of a point is called *acyclic*.

**Definition 16.2.9** (Simplicially connected). Let K be a simplicial complex. Then we call K simplicially connected iff for all pairs of 0-simplices  $v_0, v_n \in K$ , there exists a sequence of 0-simplices  $\{v_i\}_{i \in [n]}$  such that for all  $i \in [n]$  (with  $i \neq n$ ),  $\{v_i v_{i+1}\}$  is a 1-simplex in K. Note, this corresponds exactly to K being connected as a graph, where the 0-simplices represent vertices, and the 1-simplices represent edges.

**16.8.** If K is simplicially connected, then  $H_0(K) \cong \mathbb{Z}/2\mathbb{Z}$ . If K has r simplicially connected components, then

 $\mathsf{H}_0(K) \cong \prod_{i=1}^r \mathbb{Z}/2\mathbb{Z}$ 

Solution.

(a) Suppose K is simplicially connected. We want to show  $H_0(K) \cong \mathbb{Z}/2\mathbb{Z}$ . First, observe that  $Z_0(K) \cong C_0(K)$  (every 0 simplex has trivial boundary). By properties of module homomorphisms, for all  $\sigma \in B_0(K)$ ,  $\sigma$  is a basis element of  $B_0(K)$  iff  $\exists \tau \in C_1(K)$  such that  $\tau$  is a basis element of  $C_1(K)$ , and  $\partial_1(\tau) = \sigma$ . Thus,  $B_0(K)$  is spanned by  $\{\{\{v_i\} + \{v_j\}\} \mid \{v_iv_j\} \in K\}$ . It follows that  $B_0(K)$  contains exactly those elements of  $C_0(K)$  with an even number of vertices.<sup>2</sup>

It follows that  $H_0(K) = Z_0(K)/B_0(K) \cong \mathbb{Z}/2\mathbb{Z}$  (any 0-chain has either an even or odd number of vertices).

(b) This follows by applying the above argument to each of the connected components.

**16.9.** Let K be a triangulation of a 3-dimensional ball that consists of a 3-simplex together with its faces. Compute  $H_n(K)$  for each n.

Solution.

<sup>&</sup>lt;sup>2</sup>Since  $\mathsf{B}_0(K)$  is generated by pairs.



Figure 16.7: 3-simplex and its basis faces. Note the 1, 4, 6, 4 relationship. Gotta love Pascal's 2-simplex!

- (0) K is connected, so  $\mathsf{H}_0(K) \cong \mathbb{Z}/\mathbb{Z}2$ .
- (1) Elements of  $Z_1(K)$  are all just closed loops (linear combinations of the  $\partial_2(A_i)$ ). But elements of  $B_1(K)$  are also just linear combinations of the  $\partial_2(A_i)$ . Hence  $H_1(K) \cong \mathbf{0}$ .
- (2)  $\mathsf{Z}_2(K) = \{\mathbf{0}, A_1 + A_2 + A_3 + A_4\} = B_2(K) \cong \mathbb{Z}/2\mathbb{Z}, \text{ so } \mathsf{H}_2(K) \cong \mathbf{0}.$
- (3)  $H_3(K) \cong \mathbf{0}$ .

It follows that the 3-simplex with all its faces is acyclic, which makes sense, since the

underlying space is homeomorphic to the 3-ball, and the 3-ball is homeomorphic to a point.  $\hfill\blacksquare$ 

**16.10.** Let K be a triangulation of a 2-sphere that consists of the proper faces of a 3-simplex. Compute  $H_n(K)$  for each n.

Solution. Proceed as before for k = 0, 1. For k = 2, note  $\mathsf{B}_2(K) \cong \mathbf{0}$ . Hence,  $\mathsf{H}_2(K) \cong \mathbb{Z}/2\mathbb{Z}$ .

**Definition 16.2.10** (Seeing a simplex). Let K be a simplicial complex with  $|K| \subset \mathbb{R}^n$ . A point  $x \notin K$  can see K if any ray from x intersects |K| at most once (as seen in the following diagram).



Figure 16.8: Simplices being seen

**Remark.** Note, that when there are multiple k-simplices in K, the picture might not be quite as simple.

**Remark.** As far as I can tell, a point x sees K iff x is in orthogonal complement of the k-hyperplane containing K. Not sure if this is actually correct though?

**Definition 16.2.11** (Cone of x over  $\sigma$ ). Let K be a finite complex and x a point that sees K. If  $\sigma = \{v_0 \cdots v_k\}$  is a simplex of K, define the *cone* of x over  $\sigma$  to be the simplex

$$\mathsf{Cone}_x(\sigma) = \{xv_0\cdots v_k\}.$$

Last Updated April 19, 2019

 $21 \ \mathrm{of} \ 54$ 



Figure 16.9: Some cones

**Definition 16.2.12** (Cone over K). Define x \* K, the *cone over* K to be the simplicial complex comprising all simplices  $Cone_x(\sigma)$  for  $\sigma \in K$ , and all faces of such simplices.

**Remark.** Essentially, this just includes the base point and edges in fig. 16.9a and the base point and edges *and* faces in fig. 16.9b.



Figure 16.10:  $x * \sigma_2$ . Note, each of the faces is colored to indicate inclusion in the complex.

**Definition 16.2.13** (Simplicial cone operator). Define the simplicial cone operator  $Cone_x : C_n(K) \to C_{n+1}(x * K)$  by extending the definition of  $Cone_x(\sigma)$  linearly to chains.

**16.11.** For x seeing K, and  $\sigma$  a simplex of K,

 $\partial \operatorname{Cone}_x(\sigma) + \operatorname{Cone}_x(\partial \sigma) = \sigma.$ 

Solution. Let  $\sigma = \{v_0 \cdots v_k\}$ . Then

$$\partial\operatorname{Cone}_{x}(\sigma) + \operatorname{Cone}_{x}(\partial\sigma) = \left( \left\{ \widehat{x}v_{0}\cdots v_{k} \right\} + \sum_{i\in[k]} \left\{ v_{0}\cdots \widehat{v_{i}}\cdots v_{k} \right\} \right) + \operatorname{Cone}_{x} \left( \sum_{i\in[k]} \left\{ v_{0}\cdots \widehat{v_{i}}\cdots v_{k} \right\} \right)$$

Last Updated April 19, 2019

 $22 \ {\rm of} \ 54$ 

$$= \sigma + \sum_{i \in [k]} \{v_0 \cdots \widehat{v_i} \cdots v_k\} + \{v_0 \cdots \widehat{v_i} \cdots v_k\}$$
$$= \sigma + \sum_{i \in [k]} \mathbf{0}$$
$$= \sigma$$

as desired.

**16.12.** For any complex K and x seeing K, the complex x \* K is acyclic.

Solution.

16.13.

Solution.

#### 16.3 Induced Homomorphisms and Invariance

Fix two simplicial complexes K and L.

**16.14.** Let  $f: K \to L$  be a simplicial map. Carefully write out the definition of the natural induced map from *n*-chains of K to *n*-chains of L:

$$f_{\#n}: \mathsf{C}_n(K) \to \mathsf{C}_n(L)$$

and show that it is a homomorphism.

Solution. We define  $f_{\#}$  by its action on basis elements, then apply linear extension. Let  $\sigma = \{v_0 \cdots v_n\} \in \mathsf{C}_n(K)$  be a basis element. Then define

$$f_{\#n}(\sigma) = \begin{cases} \mathbf{0} & \text{if } f(\sigma) \text{ is not a } n\text{-simplex, and} \\ f(\sigma) & \text{otherwise} \end{cases}$$

We now apply linear extension. That is, for all  $\tau = \sum_{i \in I} \left\{ v_0^{(i)} \cdots v_n^{(i)} \right\} \in \mathsf{C}_n(K)$ , define

$$f_{\#n}(\tau) = f_{\#n}\left(\sum_{i \in I} \sigma_i\right)$$
$$= \sum_{i \in I} f_{\#n}(\sigma_i)$$

we want to show this is a homomorphism.

(1) Let  $\sigma_1, \sigma_2 \in \mathsf{C}_n(K)$  be arbitrary. Then

$$f_{\#n}(\sigma_1 + \sigma_2) = f_{\#n}\left(\sum_{k \in I \cup J} \left\{ v_0^{(k)} \cdots v_n^{(k)} \right\} \right)$$
$$= f_{\#n}\left(\sum_{i \in I} \left\{ v_0^{(i)} \cdots v_n^{(i)} \right\} + \sum_{j \in J} \left\{ v_0^{(j)} \cdots v_n^{(j)} \right\} \right)$$

Last Updated April 19, 2019

 $23~{\rm of}~54$ 

$$= f_{\#n}\left(\sum_{i\in I} \left\{ v_0^{(i)} \cdots v_n^{(i)} \right\} \right) + f_{\#n}\left(\sum_{j\in J} \left\{ v_0^{(j)} \cdots v_n^{(j)} \right\} \right)$$
$$= f_{\#n}(\sigma_1) + f_{\#n}(\sigma_2)$$

(2) We want to show  $f_{\#n}(\mathbf{0}) = \mathbf{0}$ . Note, for any  $\sigma \in \mathsf{C}_n(K)$ ,  $\mathbf{0} = \sigma + \sigma$ . By linearity,  $f_{\#n}(\mathbf{0}) = f_{\#n}(\sigma + \sigma) = f_{\#n}(\sigma) + f_{\#n}(\sigma) = \mathbf{0}$  as well, as desired.

thus  $f_{\#n}$  is a homomorphism.

The map  $f_{\#n}$  is called the *induced chain map*. The next exercise contains an important technicality about the induced chain map in the case where the image of an *n*-simplex is an (n-1)-simplex.

**16.15.** If the simplicial map  $f: K \to L$  maps an *n*-simplex  $\sigma$  to an (n-1)-simplex  $\tau$ , what is  $f_{\#n}(\sigma)$ ?

Solution. By the definition given above,  $f_{\#n}(\sigma) = \mathbf{0}$ .

**16.16.** Let  $f: K \to L$  be a simplicial map, and let  $f_{\#}$  be the induced map  $f_{\#}: \mathsf{C}_n(K) \to \mathsf{C}_n(L)$ . Then for any chain  $c \in \mathsf{C}_n(K)$ ,

$$\partial(f_{\#}(c)) = f_{\#}(\partial(c))$$

In other "words," we have the following commutative diagram

#### Figure 16.11: Commutative diagram

Solution. Let  $c \in C_n(K)$ . Express c as a sum of basis elements  $\{\sigma\}_{i \in I}$ . Let  $\{\sigma_i\}_{i \in I'}$  be those  $\sigma_i$  for which  $f_{\#}(\sigma_i) \neq \mathbf{0}$ . Then

$$\begin{aligned} \partial_n(f_{\#n}(c)) &= \partial_n \left( \sum_{i \in I'} f_{\#n}(\sigma_i) \right) \\ &= \partial_n \left( \sum_{i \in I'} f(\sigma_i) \right) \\ &= \partial_n \left( \sum_{i \in I'} \left\{ f\left( v_0^{(i)} \right) \cdots f\left( v_n^{(i)} \right) \right\} \right) \\ &= \sum_{i \in I'} \sum_{j=1}^n \left\{ f\left( v_0^{(i)} \right) \cdots \widehat{f\left( v_j^{(i)} \right)} \cdots f\left( v_n^{(i)} \right) \right\} \\ &= \sum_{i \in I'} \sum_{j=1}^n f_{\#n-1} \left( \left\{ v_0^{(i)} \cdots \widehat{v_j^{(i)}} \cdots v_n^{(i)} \right\} \right) \end{aligned}$$

Last Updated April 19, 2019

24 of 54

$$= \sum_{i \in I'} f_{\#n-1}(\partial(\sigma_i))$$
$$= f_{\#n-1}\left(\partial_n\left(\sum_{i \in I'} \sigma_i\right)\right)$$
$$= f_{\#n-1}(\partial_n(c))$$

as desired.

**Definition 16.3.1** (Induced Homomorphism). Let  $f : K \to L$  be a simplicial map. The *induced* homomorphism  $f_* : H_n(K) \to H_n(L)$  is defined by  $f_*([z]) = [f_{\#}(z)]$  (where the square brackets indicate an equivalence class).

**16.17.** Let  $f: K \to L$  be a simplicial map. Then the induced homomorphism  $f_*: H_n(K) \to H_n(L)$  is a well-defined homomorphism.

Solution. That  $f_*$  is a homomorphism follows directly from the definition.

- (1) That  $f_*([\mathbf{0}]) = [\mathbf{0}]$  follows by the definition of  $f_{\#}$ .
- (2) Similarly for  $f_*(\sigma + \tau) = f_*(\sigma) + f_*(\tau)$ .

We now show that  $f_*$  is well-defined. Let  $[\sigma] \in H_n(K)$  and  $[\tau] \in H_n(K)$  with  $[\sigma] = [\tau]$ . Then  $\exists \rho \in B_n(K)$  s.t.  $\sigma = \tau + \rho$ . Observe that

$$f_*([\sigma]) = [f_{\#}(\sigma)] \\= [f_{\#}(\tau + \rho)] \\= [f_{\#}(\tau) + f_{\#}(\rho)] \\= [f_{\#}(\tau)] + [f_{\#}(\rho)] \\= [f_{\#}(\tau)] + [\mathbf{0}] \\= [f_{\#}(\tau)],$$

as desired.

**16.18.** Let K be a complex comprising the proper faces of a hexagon: six edges and six vertices  $v_0, \ldots, v_5$ . Let L be the complex comprising the proper faces of a triangle: three edges and three vertices  $w_0, w_1, w_2$ . Let f be a simplicial map that sends  $v_i$  to  $w_{i \mod 3}$ . Compute the homology groups of K and L and describe the simplicial map f and the induced homomorphism  $f_*$ .

Solution.

- (1) We compute the homology groups of K. Observe,  $H_2(K) \cong \{0\}$  (since  $Z_n(K)$  is trivial).  $H_1(K) \cong \mathbb{Z}/2\mathbb{Z}$  (since  $Z_1(K) \cong \mathbb{Z}/2\mathbb{Z}$ , as we either have the whole hexagon or we don't, and  $B_1(K) \cong \{0\}$ ). Finally, by theorem 16.8, we have  $H_0(K) \cong \mathbb{Z}/2\mathbb{Z}$ .
- (2) The homology groups of L are the same.
- (3) The map f folds the circle |K| onto itself
- (4)  $f_*$  is an isomorphism.

**Definition 16.3.2** ( $\lambda$ -map). Let K be a simplicial complex. Let  $\lambda : \operatorname{sd} K \to K$  be defined as follows: for any vertex  $v \in \operatorname{sd} K$ , there exists  $\sigma \in K$  such that v is the barycenter of  $\sigma$ . Then let

$$\lambda(v) = v_{\sigma}$$

where  $v_{\sigma}$  is a vertex in  $\sigma$ .

**Definition 16.3.3**  $(\lambda_*)$ . Let  $\lambda_* : H_n(\mathsf{sd} K) \to H_n(K)$  be defined by linear extension of  $\lambda$  to simplices. Since  $\lambda$  is a well-defined simplicial map,  $\lambda_*$  is a well-defined homomorphism (theorem 16.17).

**16.19.** Suggest a homomorphism from  $C_n(K) \to C_n(\operatorname{sd} K)$  that commutes with  $\partial$ . Could its induced homomorphism on homology be an inverse for  $\lambda_*$ ?

Solution. Consdier  $f : \mathsf{C}_n(K) \to \mathsf{C}_n(\mathsf{sd}\, K)$  defined by

 $f(\sigma) = \text{sum of maximal } n \text{-simplices in } \mathsf{sd} \sigma$ 

We give this a name.

**Definition 16.3.4** (Subdivision operator). Define the subdivision operator  $SD : C_n(K) \to C_n(sd K)$  by first defining SD on a simplex:

$$\mathsf{SD}(\{v_0\cdots v_n\}) = \sum_{\pi\in\mathcal{S}_{n+1}} \{b_0^{\pi}\cdots b_n^{\pi}\}$$

where  $S_{n+1}$  is the symmetric group, and  $b_k^{\pi}$  is the barycenter of the face  $\{v_{\pi(0)}\cdots v_{\pi(k)}\}$ .

**Note.** I'm pretty sure this gets us all the maximal simplices, as we want. To verify it, I think we proceed as follows: let  $\sigma \in \operatorname{sd} K$  be an arbitrary *n*-simplex. Then one of the vertices in  $\sigma$  is a vertex in K (def. of maximal simplex). Take  $\pi$  such that  $v_{\pi(0)}$  is this maximal vertex. Also restrict  $\pi$  such that  $v_{\pi(1)}$  is the vertex such that the barycenter of  $\{v_{\pi(0)}v_{\pi(1)}\}$  is in  $\sigma$  (again, since  $\sigma$  is maximal, I think this works). Continue this.

I think also that one can show this necessitates the resulting simplices be disjoint?

**16.20.** The subdivision operator commutes with the boundary operator, that is, if c is a chain in K, then  $SD(\partial c) = \partial SD(c)$ .

Solution. We show the result for a simplex. For intuition, observe the following diagram in the case c is a 2-simplex:



Figure 16.12: SD(c)

under  $\partial$ , only the solid lines are not annihilated. In generality,

$$\begin{split} \partial \operatorname{SD}(c) &= \partial \sum_{\pi \in \mathcal{S}_{n+1}} \{ b_0^{\pi} \cdots b_n^{\pi} \} \\ &= \sum_{\pi \in \mathcal{S}_{n+1}} \sum_{j=0}^n \left\{ b_0^{\pi} \cdots \widehat{b_j^{\pi}} \cdots b_n^{\pi} \right\} \\ &= \sum_{j=0}^n \sum_{\pi \in \mathcal{S}_{n+1}} \left\{ b_0^{\pi} \cdots \widehat{b_j^{\pi}} \cdots b_n^{\pi} \right\} \\ &= \sum_{j=0}^{n-1} \sum_{\pi' \in \mathcal{S}_n} \left\{ b_0^{\pi'} \cdots b_{n-1}^{\pi'} \right\} \\ &= \operatorname{SD}(\partial c) \end{split}$$

as desired.<sup>3</sup>

**Note.** I think that it's getting a little tricky here to see which concepts are the "important parts." Maybe let's shift to trying http://www.indiana.edu/ lniat/m621notessecondedition.pdf

## 16.4 The Mayer-Vietoris Theorem

**Definition 16.4.1** (Subcomplex). If K is a simplicial complex, a *subcomplex* is a simplicial complex L such that  $L \subset K$ .

**Note.** The thing to note here is that if we choose some simplex to be in our subcomplex, we must bring all its faces with us as well.

**16.31.** If K is a finite simplicial complex, verify that the intersection of two subcomplexes of K is a subcomplex.

 $<sup>{}^{3}\</sup>pi'$  is the permutation given by  $\pi^{-1}(j_0) = \pi(j_0)$ , where  $j_0$  is the deleted vertex.

Solution. Let L, M be subcomplexes of K. Then for all  $\sigma \in L \cap M$ ,  $\sigma \in L$ ,  $\sigma \in M$ , hence for all faces  $\sigma' \subset \sigma$ , we have  $\sigma' \in L$ ,  $\sigma' \in M$ , and thus  $\sigma' \in L \cap M$ . Hence  $L \cap M$  is a simplicial complex.

The disjointness condition follows similarly.

We'll now examine cases where we have two subcomplexes A, B of a simplicial complex K. We want to look at relationships between cycles in  $A, B, A \cap B$ , and K.



Figure 16.13: An example of such A, B

**16.32.** Note that a cycle in  $A \cap B$  is still a cycle in A, B, and K. Then answer:

- (a) Can a trivial cycle in  $A \cap B$  be non-trivial in A?
- (b) Can a non-trivial cycle in  $A \cap B$  be trivial in A?
- (c) Can a non-trivial cycle in  $A \cap B$  that's also non-trivial in A and in B be trivial in K?

Solution. Let  $\sigma \in A \cap B$ . I'll assume this is asking us to just consider just the inclusion map applied to  $\sigma$ 

- (a) Nope. Including into A won't change  $\sigma$  at all.
- (b) No?
- (c) No?

**Definition 16.4.2** ("Intersection" map). Let A, B be subcomplexes of a simplicial complex K. Define the homomorphisms  $\pi_A : \mathsf{Z}_k(K) \to \mathsf{Z}_k(A), \pi_B : \mathsf{Z}_k(K) \to \mathsf{Z}_k(B)$  as follows:

$$\pi_A(\sigma) = \begin{cases} \sigma & \text{if } \sigma \in \mathsf{Z}_k(A) \\ \mathbf{0} & \text{otherwise} \end{cases} \qquad \pi_B(\sigma) = \begin{cases} \sigma & \text{if } \sigma \in \mathsf{Z}_k(B) \\ \mathbf{0} & \text{otherwise} \end{cases}$$

Observe that the following diagram commutes, and that  $\pi_A$ ,  $\pi_B$  are idempotent:<sup>4</sup>

<sup>&</sup>lt;sup>4</sup>Ok, technically dom  $\pi_A = Z_k(K) \neq Z_k(B)$ , but you could throw an inclusion map in there if you so pleased.

$$\begin{aligned}
 Z_k(K) & \xrightarrow{\pi_A} Z_k(A) \\
 \pi_B & \swarrow \\
 Z_k(B) & \xrightarrow{\pi_A} Z_k(A \cap B)
 \end{aligned}$$

Figure 16.14:  $\pi_A \circ \pi_B = \pi_B \circ \pi_A$ 

Let K be a finite simplicial complex and A and B be subcomplexes such that 16.33.  $K = A \cup B$ . If  $\alpha$ ,  $\beta$  are k-cycles in A and B respectively, and if  $\alpha \sim_{\mathbb{Z}/2\mathbb{Z}} \beta$  in K, then there is a k-cycle c in  $A \cap B$  such that  $\alpha \sim_{\mathbb{Z}/2\mathbb{Z}} c$  in A and  $\beta \sim_{\mathbb{Z}/2\mathbb{Z}} c$  in B.

Solution. The question can be rephrased as

Let  $\alpha \in \mathsf{Z}_k(A)$ , and  $\beta \in \mathsf{Z}_k(B)$ . Suppose that

$$[\alpha]_K = [\beta]_K$$

Then there exists  $c \in \mathsf{Z}_k(A \cap B)$  such that

$$\alpha]_A = [c]_A \qquad [\beta]_B = [c]_B$$

 $[\alpha]_A = [c]_A \qquad [\beta]_B = [c]_B$ Or, show that if  $\alpha - \beta = 0 \in \mathsf{H}_k(K)$ , then  $\exists c \in \mathsf{Z}_k(A \cap B)$  such that  $(\alpha, \beta) = (c, c) \in \mathsf{H}_k(A) \oplus \mathsf{H}_k(B)$ . This gives us maps

$$\mathsf{H}_n(K) \xrightarrow{\delta^k} \mathsf{H}_k(A \cap B) \xrightarrow{\phi^k} \mathsf{H}_k(A) \oplus \mathsf{H}_k(B)$$

by

$$[\alpha] = [\beta] \stackrel{\delta^k}{\longmapsto} [c] \stackrel{\phi^k}{\longmapsto} [(c,c)]$$

Since  $[\alpha]_K = [\beta]_K$ , there exists  $c_0 \in \mathsf{B}_k(K)$  such that  $\alpha - \beta = c_0$ . By definition of  $\mathsf{B}_n(K)$ , this implies that there exists  $\gamma \in \mathsf{C}_{k+1}(K)$  with  $c_0 = \partial \gamma$ .

**Claim:**  $c = \alpha + \partial \pi_A(\gamma)$  works.

#### **Proof of Claim:**

(a) First, we show c is a k-cycle. Note

$$\partial c = \partial \alpha + \partial^2 \pi_A(\gamma)$$
$$= \partial \alpha = \mathbf{0}$$

as desired.

(b) Now, we verify that equivalences. First, note that  $c - \alpha = \partial \pi_A(\gamma)$ , hence  $[\alpha]_A = [c]_A$ trivially. Now,

$$c - \beta = c - \alpha + \alpha - \beta \qquad (0 = \alpha - \alpha)$$
  
=  $(c - \alpha) + \alpha - \beta$  (grouping)  
=  $\partial \pi_A(\gamma) + \alpha - \beta$   $(c - \alpha = \partial \pi_A(\gamma))$ 

 $= \partial \pi_A(\gamma) + c_0 \qquad (c_0 = \alpha - \beta)$  $= \partial \pi_A(\gamma) + \partial \gamma \qquad (\partial \gamma = c_0)$  $= \partial (\pi_A \gamma + \gamma) \qquad (\partial \text{ commutes})$ 

hence  $c - \beta \in \mathsf{B}_k(K)$ . So  $[c]_B = [\beta]_B$ , as desired.

**16.34.** Let K be a finite simplicial complex and A and B be subcomplexes such that  $K = A \cup B$ . Let z be a k-cycle in K. Then there exist k-chains  $\alpha$  and  $\beta$  in A and B respectively such that:

z = α + β and
∂α = ∂β is a (n - 1)-cycle c in A ∩ B.
If z = α' + β', a sum of n-chains in A and B respectively, and c' = ∂α' = ∂β' is a (n - 1)-cycle, then c' is homologous to c in A ∩ B.

Solution.

(1) Let 
$$\alpha = \pi_A(z)$$
. Then  $\alpha \in \mathsf{C}_k(A)$ . Now, taking  $\beta = z - \alpha$ , we see

$$\pi_A(\beta) = \pi_A(z - \alpha)$$
  
=  $\pi_A(z - \pi_A(\alpha))$   
=  $\pi_A(z) - \pi_A(\pi_A(z))$   
=  $\pi_A(z) - \pi_A(z)$   
=  $\mathbf{0}$ 

hence  $\beta \in \mathsf{C}_k(B)$ .

(2) WTS 
$$\partial \alpha = \partial \beta \in \mathsf{Z}_{n-1}(A \cap B)$$
. Note,

$$\partial(\beta) = \partial(z - \alpha)$$
  
=  $\partial(z) - \partial(\alpha)$   
=  $\mathbf{0} - \partial(\alpha) = \partial(\alpha),$ 

which are (n-1)-boundaries (and hence (n-1)-cycles). Projection onto  $A \cap B$  yields the desired result.

(3) In  $A \cap B$ , c and c' are both (n-1) boundaries. Hence, they are trivially homologous.

**16.36.** Let K be a simplicial complex and A and B be subcomplexes such that  $K = A \cup B$ . Construct natural homomorphisms  $\phi, \psi, \delta$  among the groups below and show that  $\psi \circ \phi = 0$ and  $\delta \circ \psi = 0$ .

(a)  $\phi : \mathsf{H}_n(A \cap B) \to \mathsf{H}_n(A) \oplus \mathsf{H}_n(B).$ (b)  $\psi : \mathsf{H}_n(A) \oplus \mathsf{H}_n(B) \to \mathsf{H}_n(K).$ (c)  $\delta : \mathsf{H}_n(K) \to \mathsf{H}_{n-1}(A \cap B).$ 

Solution. Let

(1) 
$$\phi: \mathsf{H}_n(A \cap B) \to \mathsf{H}_n(A) \oplus \mathsf{H}_n(B)$$
 be given by

$$\phi(\sigma) = (\sigma, \sigma).$$

That this is a well-defined homomorphism follows immediately.

(2) Let  $\psi(\sigma, \tau)$  be given by

$$\psi(\alpha,\beta) = \alpha - \beta.$$

It is straightforward to verify this is a well-defined homomorphism.

(3) Let  $\sigma \in H_n(K)$ , and take  $\tau \in Z_n(K)$  a representative of  $\sigma$ . Then  $\exists c \in B_n(K)$  such that

 $\sigma+\tau=c.$ 

Then by the theorem above, there exists  $\alpha \in C_n(A), \beta \in C_n(B)$  such that

 $\alpha + \beta = c.$ 

Further,  $\partial \alpha$ ,  $\partial \beta$  is an (n-1)-cycle c' in  $A \cap B$ .

Let  $\delta(\sigma)$  be given by

$$\delta(\sigma) = \partial(\pi_A \sigma).$$

That this is a well-defined homomorphism follows, since all the maps included are linear.

Now, we show the composition things.

(a)

$$\psi \circ \phi(\alpha) = \psi(\alpha, \alpha)$$
$$= \alpha - \alpha$$
$$= \mathbf{0}$$

as desired.

(b) Let  $\alpha \in H_n(A)$ ,  $\beta \in H_n(B)$ . Then by the theorem above, there exists

$$\delta(\psi(\alpha,\beta)) = \partial(\pi_A(\alpha-\beta))$$
$$= \partial(\alpha - \pi_A(\beta))$$



Figure 16.15: Example of A, B with  $\alpha, \beta$ .

**Definition 16.4.3** (Exact sequence). Given a (finite or infinite) sequence of groups and homomorphisms:

$$\cdots \to G_{i-1} \xrightarrow{\phi_{i-1}} G_i \xrightarrow{\phi_i} G_{i+1} \to \cdots$$

the sequence is **exact at**  $G_i$  if and only if  $\operatorname{im} \phi_{i-1} = \ker \phi_i$ . The sequence is called an **exact** sequence if and only if it is exact at each group (except at the first and last groups if they exist).

**16.37.** Let K be a finite simplicial complex and A and B be subcomplexes such that  $K = A \cup B$ . The sequence

$$\cdots \to \mathsf{H}_n(A \cap B) \to \mathsf{H}_n(A) \oplus \mathsf{H}_n(B) \to \mathsf{H}_n(K) \to \mathsf{H}_{n-1}(A \cap B) \to \dots$$

using the homomorphisms  $\phi, \psi, \delta$  above, is exact.

(a)

16.39.	Compute the $\mathbb{Z}$	/2Z-homology	groups for	each complex	K below
--------	--------------------------	--------------	------------	--------------	---------

- (a) The bouquet of k circles (the union of k circles identified at a point).
- (b) A wedge of a 2-sphere and a circle (the two spaces are glued at one point).
- (c) A 2-sphere union its equatorial disk.
- (d) A double solid torus.

Solution.

(a) K is given by k triangles (together with their faces), with one common vertex. Hence

$$\mathsf{H}_1(K) = (\mathbb{Z}/2\mathbb{Z})^k \qquad \qquad \mathsf{H}_0(K) = (\mathbb{Z}/2\mathbb{Z})^k$$

(b) Observe that  $K = \partial \Delta^3 \cup \partial \Delta^2 / v_1 \sim v_2$  (where  $v_1, v_2$  are arbitrarily chosen vertices from  $\partial \Delta^3$ ,  $\partial \Delta^2$  respectively). There are just two elements of  $H_2(K)$ : namely, **0** and  $A_1 + A_2 + A_3 + A_4$ . Hence,  $H_2(K) \cong \mathbb{Z}/2\mathbb{Z}$ .

Now, since  $\partial^2 \Delta^3 = 0$ ,  $\mathsf{H}_1(\partial^2 \Delta^3)$  is trivial. Observe that  $\mathsf{H}_1(\partial \Delta^2) = \mathbb{Z}/2\mathbb{Z}$ . Now,

- (c)
- (d)

**16.40.** Compute the  $\mathbb{Z}/2\mathbb{Z}$ -homology groups of a torus using Mayer-Vietoris in two different ways (with two different decompositions).

*Solution.* By Mayer-Vietoris, independent of our particular decomposition, we will have the following exact sequence:



Figure 16.16: Mayer Vie-torus (heh)

Let A, B be two macaroni elbow shapes, overlapping on two cylindrical segments: Note



Figure 16.17: Decomposition 1: A below, B above

that for k > 2,  $\mathbb{T}^2$  has no k-cycles, and hence

$$\mathsf{H}_k(A \cap B) \cong \mathsf{H}_k(A) \oplus \mathsf{H}_k(B) \cong \mathsf{H}_k(\mathbb{T}^2) = \{\mathbf{0}\}.$$

by exactness of the Mayer-Vietoris sequence,  $\operatorname{im} \delta_3 = \ker \phi_2 = \{\mathbf{0}\}$ . Hence,  $\phi_2$  is one-to-one. We will use this later.

Calculating the n = 1, 0 homology groups is easy. By Theorem 16.8,  $\mathsf{H}_0(A \cap B) \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ ,  $\mathsf{H}_0(A) \oplus \mathsf{H}_0(B) \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ , and  $\mathsf{H}_0(\mathbb{T}^2) \cong \mathbb{Z}/2\mathbb{Z}$ .

Now, observe that each of  $H_2(A \cap B)$ ,  $H_2(A)$ , and  $H_2(B)$  are trivial (A and B bound no volume). Hence  $\delta_2$  is injective. Furthermore, since  $A \cap B$  is a disjoint union of two cylinders,  $H_1(A \cap B) \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ , and  $H_1(A) \cong H_1(B) \cong \mathbb{Z}/2\mathbb{Z}$ . This gives



Figure 16.18: Summary of results so far

Finally, we apply exactness. Since dom  $\psi_2 = \{\mathbf{0}\}$ , im  $\psi_2 = \ker \delta_2 = \{\mathbf{0}\}$ , hence  $\delta_2$  is 1-1. Now, note that  $\ker \psi_1 = \{(0,0), (1,1)\} = \operatorname{im} \phi_1$ . Hence,  $\ker \phi_1 \cong \mathbb{Z}/2\mathbb{Z} \cong \operatorname{im} \delta_2$ . It follows that  $\mathsf{H}_2(K) \cong \mathbb{Z}/2\mathbb{Z}$ .



Figure 16.19: Final diagram

we now take the second decomposition. For k > 2, all the results remain the same. Now,



Figure 16.20: Second Decomposition

note that  $A \cap B$  is given by two concentric cylinders. Since this is still the disjoint union of two cylinders, for k = 0, 1, 2 the homology groups  $H_k(A \cap B)$  and  $H_k(A) \oplus H_k(B)$ remain the same. Thus, an identical argument to that above shows that  $H_2(\mathbb{T}^2)$ ,  $H_1(\mathbb{T}^2)$ are likewise identical.

**FS1.** Suppose that K is a triangulation of two copies of  $\mathbb{S}^2$ , identified at a copy of  $\mathbb{S}^1$ . Let  $A \cong \mathbb{S}^2 \cong B$ . Use Mayer-Vietoris to compute  $H_n(K)$ .

Solution. As before, all the  $H_k$  are trivial for k > 2. For k = 0, we count the number of connected components. This yields

- $H_0(K) \cong \mathbb{Z}/2\mathbb{Z}$
- $\mathsf{H}_0(A) \oplus \mathsf{H}_0(B) \cong (\mathbb{Z}/2\mathbb{Z})^2$
- $\mathsf{H}_0(A \cap B) \cong \mathbb{Z}/2\mathbb{Z}$

For k = 1,

- $\mathsf{H}_1(A \cap B) \cong \mathbb{Z}/2\mathbb{Z}$
- $\mathsf{H}_1(A) \oplus \mathsf{H}_1(B) \cong (\mathbb{Z}/2\mathbb{Z})^2$
- •

16.41. Use the Mayer-Vietoris Theorem to compute  $H_n(M)$  for every compact, triangulated 2-manifold M. What compact, triangulated 2-manifolds are not distinguished from one another by  $\mathbb{Z}/2\mathbb{Z}$ -homology? What does  $H_2(M)$  tell you?

Solution. Let M be a compact triangulated 2-manifold, and call the triangulation K. Let A, B be subcomplexes of K such that  $A \cup B = K$ . Then for all k > 2, each of  $H_k(M)$ ,

 $\mathsf{H}_k(A), \mathsf{H}_k(B)$ , and  $\mathsf{H}_k(A) \oplus \mathsf{H}_k(B)$  are isomorphic to  $\{\mathbf{0}\}$ .

Suppose M can be expressed as the following connected sum:

**16.42.** Let  $p, q \in \mathbb{Z}$  be relatively prime. Calculate  $\mathsf{H}_n(L(p,q))$ , the homology of the lens space L(p,q).

## 18. Simplicial Z-Homology: Getting Oriented

#### **18.1** Orientation and Z-Homology

**Note.** We used  $H_n(K)$  to denote the  $\mathbb{Z}/2\mathbb{Z}$ -homology gorups of a complex K. This was to simplify notation. In general, the notation is more like  $H_n(K;G)$ , where G is the group of coefficients for our module. When  $G = \mathbb{Z}$ , we drop the group and write  $H_n(K)$ .

**Definition 18.1.1** (Edge orientation). For an edge  $\{vw\}$ , the two orientation classes correspond to two orderings of the vertices v and w, and are denoted [vw] and [wv]. It is customary to think of the oriented edge [vw] as an edge with an arrow pointing from v to w. We set [vw] = -[wv].



Figure 18.1: [vw] and [wv]

**Definition 18.1.2** (Triangle orientation). For a triangle  $\{uvw\}$  with vertices u, v, and w, the two orientation classes correspond geometrically to clockwise or counterclockwise orderings of the vertices when viewed a long a fixed normal.

**Definition 18.1.3** (Oriented Simplex). Let  $\{v_0 \cdots v_k\}$  be a k-simplex, and let  $\pi \in S_n$ . Then

$$[v_0\cdots v_k] = [v_{\pi(0)}\cdots v_{\pi(k)}]$$

iff  $\pi \in A_n$ . If  $\pi \notin A_n$ , we have

$$[v_0\cdots v_k] = -[v_{\pi(0)}\cdots v_{\pi(k)}]$$

An n-simplex with a chosen orientation is called a *Oriented simplex*. The boundary of a 0-simplex is defined to be 0.

**Definition 18.1.4** (*n*-chain group). The *n*-chain group of K is the free abelian group of oriented K-simplices under the equivalence relation above.

**Definition 18.1.5** (Boundary map). For  $n \ge 1$ , the boundary of an oriented n-simplex  $\sigma = [v_0 \cdots v_n]$  is

$$\partial(\sigma) = \sum_{i=0}^{n} (-1)^{i} [v_0 \cdots \widehat{v_i} \cdots v_n]$$

**18.3.** For any *n*-simplex  $\sigma$ ,

 $\partial(-\sigma) = -\partial(\sigma)$ 

Solution. Let  $\sigma = [v_0 \cdots v_n]$ . Since (0 1) is odd, then

$$[v_0\cdots v_n] = -[v_1v_0\cdots v_n]$$

hence

$$\begin{aligned} \partial(-\sigma) &= \partial([v_1v_0\cdots v_n]) \\ &= \sum_{i=0}^n (-1)^i [v_0\cdots \widehat{v_i}\cdots v_n] \\ &= [v_0v_2\cdots v_n] - [v_1v_2\cdots v_n] + [v_1v_0v_3\cdots v_n] + \dots + (-1)^n [v_1v_0\cdots v_{n-1}] \\ &= -[v_1v_2\cdots v_n] + [v_1v_2\cdots v_n] - [v_0v_1v_3\cdots v_n] + \dots + (-1)^{n+1} [v_0v_1\cdots v_{n-1}] \\ &= \sum_{i=0}^n (-1)^{i+1} [v_0\cdots \widehat{v_i}\cdots v_k] \\ &= -\partial(\sigma) \end{aligned}$$

as desired.

**Definition 18.1.6** (Boundary of an *n*-chain). The boundary of an *n*-chain is an (n-1)-chain given by

$$\partial \left( \sum_{i=1}^{k} c_i \sigma_i \right) = \sum_{i=1}^{k} c_i \partial(\sigma_i)$$

Thus, the boundary is a homomorphism

$$\partial_n : \mathsf{C}_n(K) \to \mathsf{C}_{n-1}(K)$$

**18.4.** For all  $n \ge 0$ ,

$$\partial_n \circ \partial_{n+1} = 0.$$

Solution. Let  $\sigma \in \mathsf{C}_{n+1}(K)$  be given by

$$\sigma = \sum_{i=0}^{k} c_i \sigma_i.$$

Then

$$\partial_n \circ \partial_{n+1}(\sigma) = \partial_n \left( \sum_{i=0}^k c_i \partial_{n+1}(\sigma_i) \right)$$
$$= \left( \sum_{i=0}^k c_i \partial_n \circ \partial_{n+1}(\sigma_i) \right)$$

hence it suffices to show that the claim holds on one of the  $\sigma_i.$  Note,

$$(\partial_n \circ \partial_{n+1})(\sigma_i) = \partial_n \left( \sum_{j=0}^{n+1} (-1)^j \left[ v_0^{(i)} \cdots \widehat{v_j^{(i)}} \cdots v_{n+1}^{(i)} \right] \right)$$
$$= \sum_{j=0}^{n+1} \sum_{1 \le \ell \ne j \le n+1}^n (-1)^{j+\ell} \left[ v_0^{(i)} \cdots \widehat{v_j^{(i)}} \cdots \widehat{v_\ell^{(i)}} \cdots v_{n+1} \right]$$

conceivably, doing all the algebra out works.

Last Updated April 19, 2019

**18.7.** For a finite simplicial complex K,  $H_n(K)$  is a finitely generated abelian group.

Solution. Let K be a finite simplicial complex. Then by definition,  $C_n(K)$  is finitely generated (by the two orientations of each of the *n*-simplices of K). Hence,

$$\mathsf{H}_n(K) = \mathsf{C}_n(K) / \mathsf{B}_n(K)$$

is finitely generated as well.

**18.8.** If K is simplicially connected, then  $H_0(K) \cong Z$ . If K has r connected components, then  $H_0(K)$  is a free abelian group of rank r.

Solution. Recall that K is simplicially connected iff for all pairs of vertices  $v_0, v_k$ , there exists a sequence of 0-simplices  $\{v_i\}_{1 \le i \le n}$  such that for all  $1 \le i \le n - 1$ ,  $\{v_i v_{i+1}\}$  is a 1-simplex in K.

Hence, for all 0-simplices  $v_i, v_k \in K$ , we have

$$v_j \sim v_k$$

by

$$v_j - v_k = \partial \left( \sum_{i=j}^{k-1} \{ v_i v_{i+1} \} \right).$$

thus  $H_0(K)$  is a  $\mathbb{Z}$ -module with 1 basis element, so  $H_0(K) \cong \mathbb{Z}$ .

Similarly, if K has r simplicially connected components, then  $H_0(K) \cong \mathbb{Z}^r$ .

**18.9.** If K is a one-point space,  $H_n(K) \cong 0$  for n > 0, and  $H_0(K) \cong \mathbb{Z}$ .

Solution. This follows directly as a corollary of the previous theorem.

#### 18.2 Relative Simplicial Homology

**Definition 18.2.1** (Relative Chain Group). Let K' be a subcomplex of a simplicial complex K. Then the chain group  $C_n(K')$  can be viewed as a subgroup of the chain group  $C_n(K)$  consisting of all chains that are zero on any simplex outside K'. Then we can define the group of relative chains of K modulo K' as the quotient group

$$C_n(K,K') = C_n(K)/C_n(K')$$

18.16. There is a boundary map

$$\partial_n : \mathsf{C}_n(K, K') \to \mathsf{C}_{n-1}(K, K')$$

such that  $\partial_n \circ \partial_{n+1} = 0$  for all  $n \ge 0$ .

Solution. Let  $\partial_n^K$  be the boundary map on  $\mathsf{C}_n(K)$ . Then define

$$\partial_n(c_n) = \begin{cases} 0 & \text{if } \partial^K(c_n) \in \mathsf{C}_{n-1}(K') \\ \partial^K_n(c_n) & \text{otherwise} \end{cases}$$

with linear extension.

Last Updated April 19, 2019

Definition 18.2.2 (Relative Homology Group). As one might expect, we define

 $\mathsf{H}_n(K,K') = \mathsf{B}_n(K,K')/\mathsf{Z}_n(K,K')$ 

**18.25.** The boundary map  $\partial : \mathsf{C}_n(K) \to \mathsf{C}_{n-1}(K')$  induces a well-defined map  $\partial : \mathsf{H}_n(K,K') \to \mathsf{H}_{n-1}(K').$ 

Solution. Consider the following diagram:



Figure 18.2: Diagram

we perform a diagram chase. Let  $z_n \in \mathsf{Z}_n(K, K') \subset \mathsf{C}_n(K, K')$ . Then  $\partial_n z_n = 0 \in \mathsf{C}_{n-1}(K, K')$ . Further, since  $\pi_n$  is surjective (by exactness), there exists  $c_n \in \mathsf{C}_n(K)$  such that  $z_n = \pi_n(c_n)$ .

By the commutativity, of the diagram,  $\partial_n \circ \pi_n(c_n) = \pi_{n-1} \circ \partial_n(c_n) = 0$ . Thus,  $\partial_n c_n \in \ker \pi_{n-1}$ , so by exactness,  $\partial_n c_n \in \operatorname{im} i_{n-1}$ . Hence, there exists  $c'_{n-1} \in \mathsf{C}_{n-1}(K')$  such that

$$i_{n-1}(c_{n-1}') = \partial_n c_n.$$

Last Updated April 19, 2019

40 of 54

We summarize the results so far in the following diagram:



Figure 18.3: Diagram

we now verify that  $c_{n-1}$  is a cycle. Note that  $\partial_n \circ \partial_{n+1}(c_n) = 0$ . Since  $i_{n-1}$  is injective,  $\partial_n \circ \partial_{n+1}(c_n) = i_{n-1}(0)$ . So by commutativity,



Figure 18.4: Diagram

**Definition 18.2.3** (Chain Complex). A chain complex C is a family  $\{C_n, \partial_n\}$  of abelian groups  $C_n$  and homomorphisms  $\partial_n : C_n \to C_{n-1}$  such that  $\partial_n \circ \partial_{n+1} = 0$  for all n. The *n*-th homology group  $H_n(\mathcal{C})$  is defined by

$$H_n(\mathcal{C}) = \ker \partial_n / \operatorname{im} \partial_{n+1}.$$

**Definition 18.2.4.** Given two chain complexes  $C = \{C_n, \partial_n\}$  and  $C' = \{C'_n, \partial'_n\}$ , a **chain map**  $\phi : C \to C'$  is a family of homomorphisms  $\phi_n : C_n \to C'_n$  such that the  $\phi_n$  commute with the boundary maps:

$$\partial'_n \circ \phi_n = \phi_{n-1} \circ \partial_n.$$



Solution. We proceed by diagram chase. We'll proceed by filling in the following diagram:



First, we show  $\gamma$  is injective. To that end, we want to show ker  $\gamma = \{0\}$ .

Let  $c \in C$ , and suppose  $c \in \ker \gamma$ . Then  $\gamma(c) = 0$ , hence  $\gamma(c) \in \ker r'$ . Thus by exactness, there exists  $a' \in A'$ ,  $b' \in B'$  such that q'(a') = b', and  $r'(b') = \gamma(c) = 0$ :



Now, since  $\alpha, \beta$  are surjective, there exists  $a \in A, b \in B$  such that  $\alpha(a) = a', \beta(b) = b'$ : Last Updated April 19, 2019 45

43 of 54



Figure 18.5: Surjectivity, indicated by the double arrowhead.

Now, note that  $s'(\gamma(c)) = s'(0) = 0$ . Since  $\delta$  is an isomorphism (and thus injective), it follows that  $\delta^{-1}(\gamma(c)) = \delta^{-1}(0) = 0$ . Finally, by commutativity of the square, we have s(c) = 0, so we can place them in corresponding boxes:



Figure 18.6: Injectivity, indicated by the hooked arrow, and commutativity.

Now,  $c \in \ker s$  implies  $c \in \operatorname{im} r$ , thus there exists  $\overline{b} \in B$  such that  $r(\overline{b}) = c$ . Let  $\overline{b}' = \beta(b)$ . Then by commutativity,

$$r'\left(\overline{b}'\right) = (r' \circ \beta)(\overline{b}) = (\gamma \circ r)(\overline{b}) = 0$$

and thus  $\overline{b}' \in \ker r'$ . By exactness, there exists  $\overline{a}' \in A'$  such that  $q'(\overline{a}') = \overline{b}'$ :

Last Updated April 19, 2019

 $44 {\rm ~of~} 54$ 



Figure 18.7

Using surjectivity of  $\alpha$ , there exists  $\overline{a} \in A$  such that  $\alpha(\overline{a}) = \overline{a}'$ . Then

$$r(\overline{b}) = r(q(\overline{a}))$$
$$= (r \circ q)(\overline{a})$$
$$= 0$$

where the last portion follows by exactness. Thus,  $\gamma$  is injective.

**Remark.** My argument was actually somewhat redundant here. I just didn't adjust it because that would have required lots of re- $T_EXing$  of diagrams.

We'll attempt a less-redundant argument for surjectivity. We'll fill in the following diagram:



Let  $c' \in C$ . Then  $d' = s'(c') \in D'$ , and t'(d') = t'(s'(c')) = 0:



## **18.3** Useful Exact Sequences



Solution. Follows as a corollary of the Zig-Zag Lemma.

## 18.4 The Degree of a Map

**Definition 18.4.1.** Let  $f : \mathbb{S}^n \to \mathbb{S}^n$  be a continuous map. Then  $f_* : H_n(\mathbb{S}^n) \to H_n(\mathbb{S}^n)$  is a homomorphism from  $\mathbb{Z}$  to itself. Hence it represent multiplication by some integer, called the *degree* of f and denoted deg f.

**18.38.** If  $f : \mathbb{S}^n \to \mathbb{S}^n$  is continuous, then deg f is well-defined. That is, it does not depend on the way in which we identify  $H_n(\mathbb{S}^n)$  with  $\mathbb{Z}$ .

 $Solution. \ ?$ 

**18.39.** Let  $f, g: \mathbb{S}^n \to \mathbb{S}^n$  be continuous maps. (a) If f and g are homotopic, they have the same degree. (b)  $\deg(f \circ g) = (\deg f) \cdot (\deg g)$ .

Solution.

(a) Since f, g are homotopic, there exists a continuous map  $F : \mathbb{S}^n \times [0, 1] \to \mathbb{S}^n$  such that  $F(\boldsymbol{x}, 0) = f(\boldsymbol{x})$ , and  $F(\boldsymbol{x}, 1) = g(\boldsymbol{x})$ .

(b)

## 18.5 Lefschetz

Definition 18.5.1 (Free Abelian Group). A group

$$G \cong \bigoplus_{i=1}^{n} \mathbb{Z}$$

is called the *free abelian group of rank* n.

**Definition 18.5.2** (Torsion Group). Let G be a group. Denote the *torsion subgroup*,  $G_{tor}$ , by

 $G_{\text{tor}} = \{ g \in G \mid \exists n \in \mathbb{N} \text{ s.t. } g^n = \epsilon \},\$ 

that is, the subgroup of all elements with finite order.

FK.	Prove that $G_{\text{tor}}$ is indeed a subgroup of $G$ .
Sole	ution. Let $g_1, g_2 \in G_{\text{tor}}$ . WTS $g_1 g_2^{-1} \in G_{\text{tor}}$ .

**Definition 18.5.3.** Let G be a finitely generated abelian group, which we can express as

 $G = G_{\text{free}} \oplus G_{\text{tor}},$ 

the direct sum of a free and a torsion group.

Forest Kobayashi

## 19. Some Homological Algebra

The big idea: algebraic topology assigns discrete algebraic invariants to topological spaces and continuous maps. Book for this section: James May's A Concise Course in Algebraic Topology

#### **19.1** Chain complexes

**Definition 19.1.1** (Chain/Cochain Complexes). Let R be a commutative ring. A *chain complex* over R is a sequence of maps of R-modules

$$\cdots \to X_{i+1} \xrightarrow{d_{i+1}} X_i \xrightarrow{d_i} X_{i-1} \to \cdots$$

such that  $d_i \circ d_{i+1} = 0$  for all *i*. We generally abbreviate  $d = d_i$ . A cochain complex over R is an analogous sequence

$$\cdots \to Y^{i-1} \xrightarrow{d^{i-1}} Y^i \xrightarrow{d^i} Y^{i-1} \to \cdots$$

with  $d^i \circ d^{i-1}$ .

We usually require chain complexes to satisfy  $X_i = 0$  for i < 0, and cochain complexes to satisfy  $Y^i = 0$  for i < 0. Without this distinction, the definitions are equivalent.

**Definition 19.1.2** (Some definitions). Elements of ker  $d_i$  are called cycles. Elements of im  $d_{i+1}$  are called boundaries. Write  $B_i(X) \subset Z_i(X) \subset X_i$  for the submodules of boundaries and dycles, and define the  $i^{\text{th}}$  homology group  $H_i(X)$  by

$$H_i(X) = Z_i(X) / B_i(X).$$

We write  $H_*(X)$  for the sequence of *R*-modules  $H_i(X)$ . We understand

Forest Kobayashi

#### 20. Rotman

The big idea: algebraic topology assigns discrete algebraic invariants to topological spaces and continuous maps. Book for this section: Joseph Rotman's A First Course in Algebraic Topology

#### 20.1 A sketch of the Brouwer Fixed Point Theorem

**R 0.1.** Every continuous function  $f: D^1 \to D^1$  has a fixed point.

Solution. We'll prove this without the techniques of analysis, so as to make the connection to the general argument slightly more obvious. Let f(-1) = a and f(1) = b.

- (1) Suppose a = -1 or b = 1, then we're done.
- (2) Else, a > -1 and b < 1. Consider the graph of f:

$$G = \left\{ (x, f(x)) \mid x \in D^1 \right\}$$

since f is continuous and  $D^1$  is connected, G is connected as well. Let

$$A = \{(x, f(x)) \mid f(x) > x\} \quad \text{and} \quad B = \{(x, f(x)) \mid f(x) < x\}.$$



Figure 20.1: G

And let  $\Delta = \{(x, x) \mid x \in [0, 1]\}$ . Note  $a \in A$ , and  $b \in B$ , so  $A \neq \emptyset \neq B$ .

Suppose  $G \cap \Delta = \emptyset$ . Then  $G = A \sqcup B$ . Note A, B are open in G, hence G is not connected, a contradiction.

**Definition 20.1.1** (retract). A subspace X of a topological space Y is a *retract* of Y if there is a continuous map  $r: Y \to Y$  with r(x) = x for all  $x \in X$ . Such a map is called a *retraction*.

Problem.

Forest Kobayashi

# 21. Appendix

# 21.1 List of Definitions

Chapter 15	Homological Prereqs 1
(Def 15.1.1)	<i>n</i> -cube
(Def 15.1.2)	<i>n</i> -ball
(Def 15.1.3)	<i>n</i> -sphere
(Def 15.1.4)	<i>n</i> -manifold
(Def 15.2.1)	Affine Independence
(Def 15.2.2)	Convex combination
(Def 15.2.3)	<i>k</i> -simplex
(Def 15.2.4)	Face and dimension
(Def 15.2.5)	Simplicial complex
(Def 15.2.6)	Underlying space
(Def 15.2.7)	Triangulable
(Def 15.3.1)	Simplicial map
(Def 15.3.2)	Simplicially homeomorphic
(Def 15.3.3)	Subdivision
(Def 15.3.4)	Piecewise linear
(Def 15.4.1)	Barvcenter
(Def 15.4.2)	First barycentric subdivision $(sd\sigma)$
(Def 15.4.3)	sd <i>K</i>
(Def 15.4.4)	<i>m</i> -th barvcentric subdivision
(Def 15.4.5)	Minimal face
(Def 15.4.6)	Star of vertex
Observes 10 /	7/07/ II 11
$(D_{\text{of}} 16.9.1)$	2/22 Homology
(Def  10.2.1)	n-chain
(Def  16.2.2)	$n$ -chain group $\dots$
(Def  16.2.3)	$\mathbb{Z}/2\mathbb{Z}$ boundary of a simplex
(Def  16.2.4)	$\mathbb{Z}/2\mathbb{Z}$ boundary of an <i>n</i> -chain $\dots \dots \dots$
(Def  16.2.5)	n-cycle and $n$ -boundary
(Def 16.2.6)	Homologous cycles
(Def 16.2.7)	$n^{\text{th}}$ Homology group
(Def 16.2.8)	Acyclic
(Def 16.2.9)	Simplicially connected
(Def 16.2.10)	Seeing a simplex
(Def 16.2.11)	Cone of x over $\sigma$
(Def 16.2.12)	Cone over $K$
$(Def \ 16.2.13)$	Simplicial cone operator
$(Def \ 16.3.1)$	Induced Homomorphism
(Def 16.3.2)	$\lambda$ -map 26
$(Def \ 16.3.3)$	$\lambda_* \dots \dots$
(Def 16.3.4)	Subdivision operator
(Def 16.4.1)	Subcomplex

(Def 16.4.2) (Def 16.4.3)	"Intersection" map	$28 \\ 32$
Chapter 18	${\mathbb Z}$ Homology	37
(Def 18.1.1)	Edge orientation	37
(Def 18.1.2)	Triangle orientation	37
(Def 18.1.3)	Oriented Simplex	37
(Def 18.1.4)	<i>n</i> -chain group	37
(Def 18.1.5)	Boundary map	37
(Def 18.1.6)	Boundary of an <i>n</i> -chain	38
(Def 18.2.1)	Relative Chain Group	39
(Def 18.2.2)	Relative Homology Group	40
(Def 18.2.3)	Chain Complex	42
(Def 18.2.4)		42
(Def 18.4.1)		46
(Def 18.5.1)	Free Abelian Group	47
(Def 18.5.2)	Torsion Group	47
(Def 18.5.2)		47
(DCI 10.0.0)		ті
Chapter 19	Some Homological Algebra	<b>49</b>
(Def 19.1.1)	Chain/Cochain Complexes	49
(Def  19.1.2)	Some definitions	49
Chapter 20	Botman	51
$(D_{of} 20 1 1)$	rotract	51
(Der 20.1.1)	1001a00	91